Original Research

The effect of Total-Body Resistance Exercise (TRX) on muscle endurance of female students

Fatemeh Jafari¹, Nafise Asgari², Mohammad hossein Askari³, Nafise Hekmatipour^{4*}, Maryam Ebrahimpour Roodposhti 5

1. Department of Nursing, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran.

2. MS student in Geriatic Nursing, Student Research Committe, Golestan University of Medical Sciences, Gorgan, Iran

3. MS student in Geriatic Nursing, Student Research Committe, Golestan University of Medical Sciences, Gorgan, Iran

4. Department of Nursing, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran.

5.MSc in Nursing, Faculty of Nursing and Midwifery, Qom University of Medical Sciences, Qom, Iran

*Corresponding Author: Nafiseh Hekmatipour, Department of Nursing, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran. E-mail: n-hekmatipour@aliabadiau.ac.ir. ORCID:0000-0003-3950-5375

Abstract:

Background:

Resistance exercise has an important role in increasing muscle strength and endurance. Increasing muscle endurance reduces fatigue and back pain, and also maintains body balance and musculoskeletal status. Therefore, this study was conducted to evaluate the effect of TRX resistance exercise on muscular endurance of female high school students.

Methods:

This experimental study was performed on 50 patients, who had been selected by simple random method and assigned to two groups of intervention and control. Intervention group received 8 sessions of TRX resistance exercise by two instructors as the researcher's assistants (2 sessions per week for 50-60 minutes per session). Control group received the school's routine training and exercise. The upper body endurance was measured by sit-up test and lower body endurance was measured by Scott test. The collected data were entered into SPSS-21 statistical software and analyzed by descriptive and inferential statistics.

Results:

Covariance test showed a significant difference in upper body muscle strength with elimination of pre-test effect (P<0.01, Eta=0.39). It also showed a significant difference in lower body muscle endurance with elimination of pre-test effect (P<0.01, Eta=0.67).

Discussion:

The results showed that TRX resistance exercise increased the muscular endurance of upper and lower torso in female students. Therefore, it is recommended to use this simple, lowcost, but effective training method in sport centers, homes and small spaces.

Keywords: TRX, Muscular Endurance, Students, Girls, Adolescents.

Submitted: 2 April 2022, Revised: 6 May 2022, Accepted: 10 June 2022

Introduction

Adolescents' physical health is considered as the main framework of a young society (1). Exercise and physical activity in adolescence play an important role in health and prevention of chronic diseases such as diabetes, heart disease, obesity, overweight, liver disorders, etc (2, 3). Regular exercise programs increase the physical and mental performance of individuals (4). Unfortunately, according to studies, more than 80% of the world's adolescents are not physically active (5). Most studies show that the rate of physical activity in adolescents is lower than the average standard defined World by Health Organization, which is 60 minutes of medium intensity physical activity per day (6, 7). However, physical activity in adolescence improves health in adulthood (8, 9). Most physical activity has a positive, immediate and long-term effect on adolescents' health (10). But today, due to factors such as changes in lifestyle, children's entertainment with television, computer games and the Internet, we see a decrease in mobility and physical activity in adolescents (11). According to studies, today's adolescents have less physical activity than their peers in the last decade (12). This inactivity in adolescents has many negative consequences on their health (13). Muscle weakness and decreased muscle endurance is one of the complications of inactivity in adolescent girls (14, 15). This causes an increase in musculoskeletal disorders, back pain and fatigue (16, 17). Weakness in the mid body muscles causes loss of balance, reduced efficiency and increased injury (18). Studies show a close relationship between muscle endurance and physical activity (19). Meanwhile, exercise plays an important role in increasing muscle endurance (20). Therefore, physical exercises increase endurance, joint mobility and balance, and

reduce stress, all of which contribute to increased quality of life (21, 22).

Total body resistance exercise (TRX) is one of the effective protocols for strengthening muscle strength (23). Studies have shown that 8 weeks of regular TRX training increases endurance, reduces weakness, and prevents muscle damage (24). TRX exercises are also performed with a special rope with two handles (8). This training method is designed to use body weight (25). Muscular activity in this training method is performed on the main and lateral muscles in all directions of movement (26, 27). In other words, TRX resistance exercise is an effective exercise method, which uses only a rope that allows exercise in a limited and small space, even in the absence of sport clubs or gyms (28). Therefore, resistance exercise is used as an effective method in creating maximum muscle strength along with muscle hypertrophy (29). The muscular resistance exercise rope is portable and can be connected in different places (26, 30), which makes this training method more cost-effective and practical that can be performed in different settings (31). Therefore, due to the convenience of this exercise method, researchers decided to conduct a study to investigate the effect of TRX resistance exercise on endurance of female students in Gorgan, Iran.

Method

This classic experimental study with two intervention and control groups was performed on 50 female high school students in Gorgan in 2021. The study environment included all female high schools in Gorgan. Inclusion criteria for this study were; being 13 to 15 years old, being female students, having no history of musculoskeletal abnormalities according to self-reported at the beginning of the study, and providing written informed consent from parents. Exclusion criteria were unwillingness to continue participating in sports sessions and being absent for more than two exercise sessions. The sample size of this study was calculated to be 50 people based on Okhli's study (2019), using G* Power sample size software with effect size of 0.72, test power of 80%, and confidence interval of 95% (32). Significant level of 0.05 was considered for all tests. The participants in this study were divided into two intervention (n = 25) and a control group (n = 25) by simple random allocation method. The multi-stage cluster method was used for sampling. First, the researcher prepared a list of female high schools in Gorgan. Then, by simple random sampling method, 3 schools were selected as the research environment. Then, after reviewing the health records and consulting with the school's sport coaches, the researcher prepared a list of students who were eligible to enter the study. It should be noted that before conducting this research, the research project was approved by the Faculty's Research Committee, while obtaining a bioethics code, registration right, and a clinical trial code. At first, the researcher presented an introductory letter to the school's officials and obtained their permission for sampling. The study objectives and method were explained to the educators, school's principals and students' Written informed parents. consent was obtained from parents and students. All participants were assured that their personal information will be kept confidential and they can withdraw from the study at any time. The study was performed blindly. The researcher had no information about the control or intervention groups as well as how to measure endurance strength before and after the intervention Randomization was also performed by school's sport coaches. The intervention was performed in a sport club under the supervision of a special coach with a bachelor's and master's degree in physical education.

Scott test was used to assess and measure muscle endurance of lower body (20), with following steps: The students stand with their back against the wall. Hips and knees are bend to a 90 degree angle. They are then asked to lift one foot 5 cm off the ground and hold as long as possible. The timing starts when the leg is raised and stops when the leg is lowered.

Sit-up test was used to measure upper body endurance. This test is used to check the strength of abdominal muscles. In this study, participants wear minimum sportswear. Then they lay on the ground with the legs being bent. Hands are placed next to the ears. Someone is holding the legs tightly. Then, they perform the test for 1 minute (16). In this study, two people observed and recorded the number of sit-ups for confirmation.

In total, 8 sessions (2 sessions per week) of TRX endurance exercise were held in the intervention group for 50 to 60 minutes each (Table 1). At the end of this study, to eliminate any discrimination between the two study groups, several resistance training sessions were held in the club and at home for samples in the control group that were willing to participate in resistance exercise sessions.

The collected data were entered into SPSS-21 statistical software. The result of Shapiro test did not show normality between the variables of this study. Data were analyzed by descriptive (table, mean, standard deviation) and inferential statistics (Independent t-test, Wilcoxon, Mann-Whitney and Covariance tests) at a significant level of 0.05.

Results

Independent t-test showed no significant difference between the intervention and control groups in terms of age (p=0.49), weight (p=0.36), height (p=0.48) and BMI (P=0.89). Wilcoxon test in the control group did not show a significant difference in upper body (P=0.16) and lower body muscles strength

(P=0.17) before and after the intervention (Table 2).

Wilcoxon test in the intervention group showed a significant difference in upper body (P=0.01) and lower body muscles strength (P<0.01) before and after the TRX exercise (Table 2). Mann-Whitney test did not show a significant difference between the intervention and control groups in terms of upper body (P=0.47) and lower body muscles strength (P=0.44) before the intervention (Table 2), but it showed a

TRX	Explanation	Involved
Exercise		muscles
Chest press	Position the body back on the center of the device. Grasp the TRX handles and angle the body. Bend your elbows to align your body with the surface of your chest. Then return the body to the starting point by opening the elbows triceps	Large chest, deltoid, and triceps
Suspended lounge (both legs)	Insert one of the legs into both handles so that the foot locks in the handles. Take a step from the center of the device. Now, gently bend the knee that bears the weight of the body. Remember, the foot should be in front of the knee. Then return the knee to its original position.	Quadriceps, hamstring, serine
Rowing, both hands	Turn your body towards the device. Hold the handles so that they are close together. Positioned so that the whole body is in a straight line. Lift the body up to chest level and then return to the starting position.	Trapezius, large dorsal, and deltoid
Scott	Grab both handles with both hands while facing the device. The distance between the hands should be a normal. Bend your knees so that they are behind your feet until your thighs are parallel to the ground, using your arms only as a support to maintain balance. Then return to the starting position.	Quadriceps, hamstring, serine
YTW	Hold both handles while standing in front of the device. The hands should cover the entire range of motion in three movements while doing a circular motion. This is a three-step movement that the body must return to the starting position after completing each step. The body slowly angles. Hands are raised in a Y-shape to the top of head. Then hands are brought down side ways completely open to form a T-shape. Finally, arms are lowered and placed next to thighs to form a W shape.	Deltoid, trapezius, and rhomboid muscles
Romanian deadlift (both legs)	Insert one foot into both handles. Take one step away from the center of the device. Bend over while the lower back is straight and the involved leg is slightly bent to hip level. Now bend the upper torso towards the ground. Finally return to the original position.	Hamstring and serine
Triceps	Position your back towards the device center. Hold both handles with the hands above the head and fully extended. The body should be in the same direction and the elbows should be level with the shoulders. Bend your elbows to a 90-degree angle and then return to the first stretched position.	Triceps
Hamstring	Lie down with your back on the floor. Insert the heel of each foot into the handles so that the back of the foot is in contact with them. Then pull the heels in a bridge-like position towards the hips and hold in the same position. Do inward and outward motion alternately.	Hamstring
Plank (flat)	While using your elbows on the floor as a support, lie on your back and insert your feet into the handle so that top of your feet are inside the	Rectus,
(flat)	insert your reet into the nancie so that top of your reet are inside the	ıransverse,

Table	1.	The	TRX	muscle	resistance	training	sessions	and	exercises ((33)	۱.
1 and	1.	тпс	IIM	muscic	I USISTAILU	u anning	505510115	anu	CACI CISCS	55	,

	handle. Now stay in this situation as much as possible.	and	side	
		muscle	25	
Side Hold	Stand sideways to the center of the device. Hold both handles at the	Side	and	
Isometric	metric level of the chest. Angle your body. Now move the body up and down		transverse	
with rope	by moving the cuffs up and down.	muscle	es	

Table 2: Comparison of muscle strength in the two groups before and after the intervention

	Group	Intervention	Control	P-value
Time				
Upper body	Before intervention	38.93 ± 4.81	37.97 ± 4.51	P = 0.47
muscles	After intervention	42.49 ± 4.19	36.08 v 4.04	P < 0.01
	P_Value	P = 0.01	P = 0.16	
Lower	Before intervention	32.76 ± 4.87	31.8 ± 3.84	P = 0.44
body	After intervention	43 ± 3.67	33.36 ± 2.17	P < 0.01
muscles	P_Value	P < 0.01	P = 0.17	

significant difference between the intervention and control groups in terms of upper body (P=0.01) and lower body muscles strength (P<0.01) after the intervention (Table 2).

Covariance test showed a significant difference in the upper body muscular strength with elimination of pretest effect (P<0.01, Eta=0.39), so that 39% of strength changes in upper body muscles could be explained by TRX resistance exercise.

Covariance test also showed a significant difference in the lower body muscular strength with elimination of pretest effect (P<0.01, Eta=0.67), so that 67% of strength changes in lower body muscles could be explained by TRX resistance exercise.

Discussion

The results of this study showed that TRX resistance exercise increased endurance in upper and lower body muscles of female students. These results are consistent with other studies.

Kiani (2021) stated that 8 weeks of TRX resistance exercise increases endurance and strengthen the muscles of mid body (24).

Ilbeigi (2014) showed that 6 weeks of TRX training increases muscle endurance and reduces fatigue and low back pain (16). Abbaszadeh (2018) revealed that regular exercise increases muscle endurance and mental health, and also reduces muscle injuries (34).

Lai (2021) in a study showed that TRX exercise increases muscle strength, physical fitness, and lower body strength (35). Seo (2021) argued that endurance training prevents the destruction of body mass and increases muscle strength and function (36).

Ansari (2020) in a study showed that 6 weeks of TRX training can increase the strength and endurance of upper and lower body muscles (26).

Mahmudieh (2017) revealed that endurance exercises such as griping robber band, increase grip strength and endurance (37). Nakhzari (2018) stated that exercises that increase endurance improve aerobic activity, reduce fatigue and enhance daily performance (38). Dehghani (2021) showed that 8 weeks of continuous endurance training increased muscle strength in the participants (23). Hovanloo (2021) revealed that resistance exercise (TRX), by increasing muscle coordination, increases the coordination of neuromuscular systems and also enhances endurance and joint stability (39). Emami (2020) in a study showed that TRX training techniques increase muscle strength and prevent fatigue in motor activities (40).

Mohammadi (2021) argued that resistance training increases muscle flexibility and endurance (41). Barbalho (2021) also showed that resistance exercise increases muscle endurance by creating muscle tension (42).

According to the results of these studies, it can be said that increasing endurance leads to muscle strength, followed by strengthening of pelvic floor and diaphragm muscles, and consequently balances the muscle function (43). This is while reducing endurance leads to decreased stability and body balance (44). Muscular strength and endurance stabilize the spine and improve motor function (45). Thus, since resistance exercise increases the muscle endurance of upper and lower torso, this simple but effective exercise method can be used to strengthen students' muscles. One of the limitations of this study was the lack of necessary facilities and physical space in schools for the implementation of intervention. In order to minimize the limitations in this study, the intervention instead of schools, was implemented in the gyms.

Conclusion

The results of this study showed that 8 weeks of resistance training increased the endurance of upper and lower body muscles in female students. Therefore, it can be said that doing resistance exercises strengthens muscle endurance. According to similar studies, increasing muscle endurance in girls helps to maintain balance, prevent musculoskeletal abnormalities, increase health, improve quality of life, and reduce fatigue and back pain. Therefore, since TRX resistance exercise is a simple and practical method that can be performed even in small spaces or at home, we recommend this exercise method to be used in schools by educators and school health officials to encourage students stay healthy and proactive.

Acknowledgement

This study was approved by Aliabad Kotoul Islamic Azad University with the ethics code: IR.IAU.CHALUS.REC.1400.057, and clinical **IRCTID:** trial registration code: IRCT20211212053363N1, which were obtained from the bioethics committee of the Islamic Azad University of Aliabad Kotoul, Golestan province. The researchers consider it necessary to thank and appreciate the efforts of schools' sport coaches and all students who helped us in conducting this study.

Conflicts of Interest

No conflict of interest was observed in this study.

Reference

- 1.Asefi AA, Amoozadeh Z. The Codification of Model of Significant Others Roles and Adolescence's Participation in Physical Activity and Sport. Research on Educational Sport. 2017;5(13):17-38.
- 2.Park JH, Jeon HS, Park HW. Effects of the Schroth exercise on idiopathic scoliosis: a meta-analysis. Eur J Phys Rehabil Med. 2018;54(3):440-9.
- 3.Rahmani Chegni S, Fathi M, Mohammad Hosseini Sarvak R. The Effect of Imposed Inactivity Due to Coronavirus Outbreak on Adolescent Body Composition and Vo2max. yums-armaghan. 2021;25(0):852-60.
- 4.Hekmati Pour N, Hojjati H. Effects of Exercise on Mental Health of Elderly. gums-hnmj. 2016;26(4):36-42.

- 5.Soleiman Ahari Z, Asadpour M, Mazar L, Nasirzadeh M. Evaluation of Physical Activity and its Effective Factors in Junior High School Female Students in Rafsanjan City: Application of Social-cognitive Theory, 2017-18. muq-journal. 2021;15(3):188-97.
- 6.Alasqah I, Mahmud I, East L, Alqarawi N, Usher K. Physical inactivity and its predictors among adolescents in Saudi Arabia: A cross-sectional comparison between cities with and without a Healthy Cities Program. Saudi Med J. 2021;42(8):886-94.
- 7.Parker K, Timperio A, Salmon J, Villanueva K, Brown H, Esteban-Cornejo I, et al. Activity-related typologies and longitudinal change in physical activity and sedentary time in children and adolescents: The UP&DOWN Study. Journal of sport and health science. 2021;10(4):447-53.
- 8.Akbarpour Beni M, Aghajani Z. Comparison of the effect of TRX and traditional resistance training on serum levels of some liver enzymes in inactive women. Journal of Sport and Exercise Physiology. 2022;15(2):20-8.
- 9.Codella R, Terruzzi I, Luzi L. Sugars, exercise and health. J Affect Disord. 2017;224:76-86.
- 10.Heradstveit O, Haugland S, Hysing M, Stormark KM, Sivertsen B, Bøe T. Physical inactivity, non-participation in sports and socioeconomic status: a large populationbased study among Norwegian adolescents. BMC Public Health. 2020;20(1):1010.
- 11.Wu XY, Han LH, Zhang JH, Luo S, Hu JW, Sun K. The influence of physical activity, sedentary behavior on healthrelated quality of life among the general population of children and adolescents: A systematic review. PLoS One. 2017;12(11):e0187668.

- 12.Moradi G, Mostafavi F, Piroozi B, Zareie B, Mahboobi M, Rasouli MA. The prevalence of physical inactivity in Iranian adolescents and the impact of economic and social inequalities on it: results of a National Study in 2018. BMC Public Health. 2020;20(1):1499.
- 13.Kumar B, Robinson R, Till S. Physical activity and health in adolescence. Clin Med (Lond). 2015;15(3):267-72.
- 14.Devries MC. Sex-based differences in endurance exercise muscle metabolism: impact on exercise and nutritional strategies to optimize health and performance in women. Exp Physiol. 2016;101(2):243-9.
- 15.Pikulska D, Kozinoga M, Janusz P, Kotwicki T. Back muscle function in adolescent girls treated with a rigid brace for idiopathic scoliosis: no impact of 6month brace wear on muscle strength or endurance. Stud Health Technol Inform. 2021;280:168-73.
- 16.Ilbeigi S, Nikbin L, Afzalpour ME. The effect of six weeks of core stability exercise on pain and trunk muscle endurance in girl students with chronic non-specific low back pain. thums-jms. 2014;2(2):5-13.
- 17.Mohammad Gholinejad P, Hojjati H, Ghorbani S. The Effect of Aerobic Exercise on Body Composition and Muscle Strength of Female Students at Elementary Schools of Ali Abad Katoul in 2018. International Journal of School Health. 2019;6(4):27-33.
- 18.Abbaszadeh A, Delavari MA. Comparing Balance, Function, Strength and Endurance of Selected Core Muscles in Male Volleyball Players with and Without a Chronic Ankle Sprain in Bandar Abbas in 2018: A Descriptive Study. RUMS_JOURNAL. 2019;18(3):251-66.
- 19.Can HB, Tuna F. Relation between endurance of deep cervical flexor muscles and physical activity level, perceived stress,

sleep quality, and smartphone addiction. Cranio. 2022;40(2):126-34.

- 20.Niazinejad N, Parnow A, Eslami R. Effect of linear and non-linear periodized programs on muscular strength and endurance in untrained adolescence girls. 2020;2018.
- 21.Zuo C, Li Q, Zhang L, Bo S. Effects of 6-Week Traditional and Functional Resistance Training on Arterial Stiffness and Muscular Strength in Healthy Young Men. Front Physiol. 2022;13:859402.
- 22.Su Y, Yao S, Zhou ZJ, Wu C, Wang IL, Lai CY. Effect of Acupuncture on Time-Dependent of Muscle Endurance in Female Elbow Joint: A Randomized Controlled Trial. Evid Based Complement Alternat Med. 2022;2022:8052256.
- 23.dehghani e, ghasemi g. Effects of eight week of Dynamic Neuromuscular Stabilization exercises on posture, strength and trunk endurance in educable mentally retarded students. Studies in Sport Medicine. 2021;13(29):229-52.
- 24.Kiani R, Fattahi H. Effects of Eight Weeks of TRX and CXWORX Exercises on Trunk Muscle Strength, Core Endurance, and Dynamic Balance of Female College Students. The Scientific Journal of Rehabilitation Medicine. 2021;10(2):186-201.
- 25.Yarahmadi Y, Hadadnejad M, Shojaedin SS. Effect of TRX Resistance Training on Functional Capacity and Lumbar Range of Motion of Middle Aged Men with Non-Specific Chronic Low Back Pain. The Scientific Journal of Rehabilitation Medicine. 2019;8(1):119-27.
- 26.Ansari Kolachahi S, Elmieh A, Talebi M. The effect of TRX exercises on serum levels of IGF-1 and cortisol and some healthrelated physical factors in active women. iau-tmuj. 2020;30(4):432-42.

- 27.Ghiyami Taklimi h, afroundeh r, bahram me, porvaghar mj, Hemmati s. The effect of 12 weeks of whole body resistance training (TRX) on testosterone and cortisol serum levels in elderly men. FEYZ. 2021;25(3):917-25.
- 28.Assar S, Gandomi F, Mozafari M, Sohaili F. The effect of Total resistance exercise vs. aquatic training on self-reported knee instability, pain, and stiffness in women with knee osteoarthritis: a randomized controlled trial. BMC Sports Sci Med Rehabil. 2020;12:27.
- 29.Lacio M, Vieira JG, Trybulski R, Campos Y, Santana D, Filho JE, et al. Effects of Resistance Training Performed with Different Loads in Untrained and Trained Male Adult Individuals on Maximal Strength and Muscle Hypertrophy: A Systematic Review. Int J Environ Res Public Health. 2021;18(21).
- 30.al-Fassih R, Fashi M, Ahmadizad S, Abuzari N. The effect of four weeks of totalbody resistance training (TRX) on muscular function and performance of young female swimmers. Journal of Sport and Exercise Physiology. 2022;15(1):21-32.
- 31.Rahimi M, Nazarali P, Alizadeh R. Pilates and TRX training methods can improve insulin resistance in overweight women by increasing an exercise-hormone, Irisin. J Diabetes Metab Disord. 2021;20(2):1455-60.
- 32.Okhli H, Hojjati H, Akhoundzadeh G. Comparing the Effect of the Corrective Exercises of America's National Academy of Sports Medicine and Pilates on the Correction of Lordosis among Female High School Students in Golestan Province in 2018. International Journal of School Health. 2019;6(4):1-6.
- 33.Janot J, Heltne T, Welles C, Riedl J, Anderson H, Howard A, et al. Effects of TRX versus traditional resistance training

programs on measures of muscular performance in adults. Journal of Fitness Research. 2013;2(2):23-38.

- 34.Abdelkader M. Effects of High intensity Interval Training Using the Elevation Training Mask on the Aerobic Capacity and Heart Rate Variability for Trained Athletes. International Journal of Sports Science and Arts. 2018;june.
- 35.Lai X, Bo L, Zhu H, Chen B, Wu Z, Du H, et al. Effects of lower limb resistance exercise on muscle strength, physical fitness, and metabolism in pre-frail elderly patients: a randomized controlled trial. BMC Geriatr. 2021;21(1):447-.
- 36.Seo MW, Jung SW, Kim SW, Lee JM, Jung HC, Song JK. Effects of 16 Weeks of Resistance Training on Muscle Quality and Muscle Growth Factors in Older Adult Women with Sarcopenia: A Randomized Controlled Trial. Int J Environ Res Public Health. 2021;18(13).
- 37.Mahmudieh Champiri B, Alizadeh MH, Akoochakian M, Emami H. The effect of 6 weeks resistance training with Theraband on upper and lower extremity muscle isometric strength in women with breast cancer. Journal of Applied Exercise Physiology. 2018;13(26):153-60.
- 38.Nakhzari Khodakheir J, Haghighi AH, Hamedinia MR. The Effects of Combined Exercise Training with Aerobic Dominant and Coenzyme Q10 Supplementation on Serum BDNF and NGF levels in Patients with Multiple Sclerosis. HBI_Journals. 2018;21(3):94-103.
- 39.hovanloo f, salarvand h, Barati AH. Effect of Six Weeks of Suspension Exercise (TRX) on the Rotator Cuf Muscles Strength and Shoulder Joint Proprioception in Girl Swimmers: Study in Prevention of shoulder impingement Syndrome. Journal for

Research in Sport Rehabilitation. 2021;8(16):101-11.

- 40.Emami A, Hadadnezhad M, Svoboda Z, Abbasi A. The effect of motion alteration on selected muscle in people who suffering from ankle disorder during landing. Jundishapur Scientific Medical Journal. 2020;19(1):53-68.
- 41.Mohammadi R, Pourrahim-e-Ghouroghchi A, Khajehlandi M. The Effect of 8 Weeks of Resistance Training with and Without Blood Flow Restriction on Serum Levels of Insulin-like Growth Factor-1 and Myostatin of Athletic Girls: A Semi-Experimental Study. RUMS_JOURNAL. 2021;20(1):53-68.
- 42.Barbalho M, Souza D, Coswig V, Abrahin O, Paoli A, Gentil P. The Effects of Resistance Exercise Selection on Muscle Size and Strength in Trained Women. Int J Sports Med. 2021;42(4):371-6.
- 43.Khodashenas E, Moradi H, Asadi Ghaleni M, Heydari E, Shams A, Enayati A, et al. The effect of selective training program on the static and dynamic balance of Deaf Children. medical journal of mashhad university of medical sciences. 2017;60(1):383-91.
- 44.Daneshjoo A, Eslami A, Mousavi Sadati SK. Effect of Core Stability Training on the Balance and FMS Scores of Adolescent Soccer Players. The Scientific Journal of Rehabilitation Medicine. 2020;9(2):61-70.
- 45.Karimizadeh Ardakani M, Mansori MH, Hamzeh shalamzari MH. Investigating the Relationship between Core Muscle Endurance with Movement Function and Postural Control in Blind People. The Scientific Journal of Rehabilitation Medicine. 2020;9(2):148-57.