## **Original Article**

# Characteristics of Non-Syndromic Supernumerary Teeth in (a Group of Turkish) Children

Yelda Kasimoglu1\*, Mine Koruyucu2, Figen Seymen3.

- 1. DDS, PhD, Istanbul University, Faculty of Dentistry, Department of Pedodontics, Istanbul, Turkey.
- 2. 2DDS, Assoc.Prof., Istanbul University, Faculty of Dentistry, Department of Pedodontics, Istanbul, Turkey.
- 3. 3DDS, Prof, Istanbul University, Faculty of Dentistry, Department of Pedodontics, Istanbul, Turkey.

\*correspondence: **Yelda Kasimoglu,** DDS, PhD, Istanbul University, Faculty of Dentistry, Department of Pedodontics, Istanbul, Turkey. Email: yelda.kasimoglu@istanbul.edu.tr

#### **Abstract:**

**Introduction**: The aim of this study was to investigate the prevalence, distribution, gender differences and characteristics of non-syndromic supernumerary (SN) teeth in children and adolescents.

**Methods**: In a 2 year cross-sectional study, a total of 16,986 patients aged between 6-14 years who visited our department between April 2014 and April 2016 were revised. 100 cases of supernumerary teeth were found. Patients with SN teeth were evaluated for demographic data (age and sex), location, morphology, classification and number of the SN teeth and related complications. The distribution rates of SN tooth types and the distribution of complications between sex and age groups were analyzed. Likelihood ratio test was used for statistical comparisons.

**Findings**: The prevalence of non-syndromic SN teeth was 0.59%. The mean age of cases included in the study was 13.88 years (65% male and 35% female). The most common type of SN tooth was conical type (50%), while the supplemental (29.6%), tuberculate (16.7%) and odontoma (3.7%) types were also found. 92% of the cases had one SN tooth. SN teeth were mostly located in the maxilla (95.6%) and were mostly unilateral (55.3%); followed by bilateral (23.7%) and midline (21.1%). There was no statistically significant difference between the distribution of SN teeth between males and females (p>0.05). At least one complication was found in 40% of the cases.

**Conclusion:** SN teeth can cause a variety of complications. Careful and early clinical and radiographic examinations in children are important in order to prevent complications of SN teeth.

**Keywords**: impacted tooth; supernumerary teeth; tooth abnormalities; dental anomaly; prevalence; pediatric patients.

### **Introduction:**

Teeth with more than normal number of teeth are called supernumerary (SN) teeth. This is a dental abnormality, also called hypoerdontia, that affects the number of teeth (another number abnormality is hypodontia/congenital tooth agenesis). Traditionally, SN teeth are defined as teeth

that originate from the odontogenic tissues of a dental lamina, consisting of more than a normal number in any type of tooth (1). It can be seen both in primary teeth and permanent teeth, and can be observed unilaterally or bilaterally in the maxilla, mandible or both. In more advanced versions, it occurs in multiple in both jaws,

and this is frequently encountered in syndromes such as Clediocranial dysplasia, (2). Gardner's syndrome,3 Incontinentia Pigmenti,4 which often accompanied by SN teeth.

Although the etiology of SN teeth has not been fully elucidated, both genetic and environmental factors play a role. There are several theories explaining the development of different SN teeth. According to the "dichotomy theory", a single tooth germ is divided into two, and two symmetrical or asymmetrical teeth are formed (5).

In addition, the type of SN tooth is divided according to the state of division of the tooth germ; when the tooth germ is divided into two equal parts, it causes the formation of two eumorphic teeth, also known as the "supplemental tooh". However, when the tooth is divided into two unequal parts, differently sized and shaped (heteromorphic) SN teeth can form. Another obvious etiological hypothesis is "atavism theory". According to this theory, while the number of teeth in our ancestors was higher in the past, the number of teeth decreased in time and its normal number decreased over time, and in some people, SN teeth can be found as an extension of the teeth we had in ancient times. Recently, SN teeth are thought to develop mostly with the "hyperactivity theory of the dental lamina" (6). Accordingly, with the induction of the hyperactivity of the dental lamina or its remnants, the dental papilla develops and then the enamel organ matures and a SN tooth is formed. In this theory, heteromorphic teeth are formed by the hyperactivity of the dental lamina residues

and eumorphic teeth from the lingual extension of an additional tooth germ (7).

SN teeth can be classified according to different parameters. In the last classification in the literature based on the morphology, position, and direction of the SN teeth; they are classified into conical, tuberculate, supplemental and odontoma according to their shape; they are also classified mesiodens, into distomolar, paramolar and parapremolar according to their location.8 They can be positioned labially, buccally or palatally in jaws; can be vertical. horizontal or inverted. treatment need may vary depending on the type of SN teeth, impaction and other factors. SN teeth can be clinically asymptomatic and may cause complications such as crowding, delayed eruption, midline diastema, dental caries, root resorption as well (9).

Hyperdontia constitutes approximately 1-3% of all dental anomalies,1 its prevalence in permanent teeth is reported to be 3.8% and the prevalence in primary teeth is 0.35-0.6% (10). The prevalence of SN teeth varies according to demographic factors, gender and race. Its prevalence in populations of Asian (11) and African (12) descent has been shown to be higher than that of the white race. SN teeth are most frequently found on the midline of the maxilla, and secondly the maxillary molar region (13).

In this study, it was aimed to investigate the prevalence of SN teeth in Turkish population and the characteristics of SN teeth in 100 cases.

#### **Methods:**

This cross-sectional study was carried out between April 2014 and April 2016 at Istanbul University Faculty of Dentistry Department of Pedodontics. The study was approved by the Istanbul University Faculty of Medicine Clinical Research Ethics Committee (No: 2013/1068).

This study included children and adolescents aged 6-14 years who applied for a dental examination on the specified dates, who had SN tooth during routine dental examination, or who had impacted SN tooth during routine radiographic examination. Patients with any syndrome, cleft/lip palate, history of jaw fracture, endodontic treatment, tooth extraction, orthodontic treatment radiographs with inadequte quality were exluded from the study. In addition, patients who had tooth loss due to dental trauma or dental caries were excluded from the study. Written informed consent was obtained from individuals who volunteered participate in the study, and intra-oral photographs were recorded within their permission.

Radiographic images was requested from patients with SN teeth during oral Panoramic radiographies examination. (Kodak 8000 Digital Panoramic Machine, Carestream Health, Inc., Rochester, NY, USA), intraoral radiographies (Kodak 2100 Intraoral Machine, Kodak, Atlanta, GA, USA) were taken with the same device. Clinical and radiographic findings were recorded by one observer (YK) for the absence or presence of SN teeth. SN teeth classified according to their were morphology, number, location and direction, and related complications were evaluated.

The families of the patients with SN tooth were also questioned in terms of the presence of SN tooth. Radiographic examination was performed for other family members who had SN tooth. Families were questioned in terms of consanguineous marriage.

Data analysis was performed using the statistical Package for the Social Sciences (SPSS) version 26 (IBM Corp, USA). Descriptive statistics of continuous and categorical variables such as gender, age, single of multiple SNs of cases were In comparing difference analyzed. complication distributions among age groups, the alternative Likelihood Ratio Test the prerequisitites was used because required for the Chi Square Independence test were not met. The significance level was set as 0.05.

## **Findings:**

This study was carried out among 100 cases who had SN teeth among 16,986 patients who applied to Istanbul University Faculty of Dentistry Department of Pedodontics. Accordingly, the prevalence of SN teeth in this patient group was found to be 0.59%.

The average age of the 100 cases included in the study was 13.88 years. 65% of the cases were boys and 35% were girls. While the most common type of SN tooth was conical type (50%); supplemental (29.6%), tuberculate (16.7%) and odontoma (3.7%) type of SN were also found. In most cases (92%), there is only one SN tooth. SN teeth were frequently localized in the maxilla (95.6%); they were mostly seen as unilateral (55.3%), then bilateral (23.7%) and midline (21.1%). As a result of clinical and

radiographic examinations, 80.6% of SN teeth were vertical, 11.1% were horizontal and 8.3% were inverted. Approximately half (50.9%) of the SN teeth were fully erupted, while the rest was partially impacted (6.4%) and impacted (42.7%). Other descriptive findings obtained in the study are given in Table 1.

The most common type of SN tooth were conical type in mixed dentition (65.5%) and permanent dentition (44.3%) period. The types of SN teeth were not differed according to the dentition (p>0.05) (Table 2).

The conical type of SN tooth were 43.2% in girls and 53.5% in boys. There was no statistically significant difference between the distribution of SN teeth according to gender groups (p>0.05) (Table 3).

At least one complication was encountered in 40% of the cases. SN teeth mostly caused delayed eruption of adjacent teeth (52.8%), followed by ectopic eruption (25%), median diastema (13.9%) and malocclusion (8.3%) (Table 1) (Fig. 1). There was no statistically significant difference between complications types between mixed and permanent dentition (p>0.05) (Table 4).

In this study, 50 cases with SN tooth and 50 families with 50 SN tooth were questioned about consanguineous marriage. The percentage of consanguineous marriage between parents of the cases with SN tooth was 76% and the families with SN tooth was 74% (Table 5).

## **Discussion and Conclusion:**

SN teeth are an uncommon developmental abnormality and can be seen in any area of the dental arch. SN teeth can be part of a genetic syndrome or can occur in healthy

non-syndromic individuals. Theories such as atavism, dichotomy of the tooth germs, excessive growth of the dental lamina have been proposed, but the etiology has not been established yet (14). It has been reported that SN teeth are more common in permanent dentition, than primary and mixed dentition. In this study, intra-oral findings of 100 nonsyndromic children with SN teeth were examined. It has been reported in the literature that the frequency of SN teeth is 0.1% to 3.8% compared to the general population (15–17). Within the scope of our study, the prevalence of non-syndromic SN teeth was found to be 0.59% in 16,986 pediatric patients. The prevalence of SN teeth was found to be higher (0.3-0.33%) than in two studies, (18,19) and lower than two other studies, (20,21) which were conducted in Turkey. The percentage of SN teeth in girls and boys was found to be similar in the study of Kara et al. (p>0.05) (18). It has been reported that SN teeth are more common in boys than girls in the literature (15,22–24).

After the tooth eruption, dental anomalies related to the crown of the teeth become detectable in the oral cavity. Studies have reported that SN teeth are detected in individuals between the ages of 5 and 70, but are most common in children between the ages of 7 and 10 (25–29). Esenlik et al.16 found that 69 of 2599 children between 6-16 years old had SN teeth with a prevalence of 2.7%. Similarly, Küchler et al.17 found the prevalence of SN teeth to be 2.3% among 1166 children examined between 6 to 12 years old. In this study, the frequency of non-syndromic SN teeth was

found to be lower than 0.59% in 16,986 children aged between 6 to 14 years.

SN teeth can be found in any area of the jaws, and the anterior maxilla has been reported to be the most affected site in the literature (16,22,30–32). In this study, it was observed that most of the SN teeth were in the maxilla (95.6%) in accordance with the literature. The percentage of maxillary SN teeth was found higher than other studies (18,33,34).

When classified according their morphology, SN teeth are divided into conical, tuberculate, supplemental and odontoma. It is stated that the most common SN tooth is of the conical type and it is generally seen among the maxillary central incisors as mesiodens. Tuberculate shaped SN teeth are mostly seen in permanent dentition and often interfere with tooth eruption. Supplemental type of SN teeth are found in both permanent and primary dentition (35). Ferres-Pedro et al (found that the most common type of SN teeth between the ages of 5-19 was conical type(22). (53.16%). Sharma et al (32). stated that the SN teeth were mostly encountered in children between the ages of 4 and 14, they were mostly seen in the central incisor region with a prevalence of 81.2% and that 30% of these SN teeth were of the conical type. According to the literature, it seems that the most common type of SN teeth in children is conical type. Similarly, the most common (50%) type of SN were found as conical type in our study.

SN teeth can be seen in the same individual as single or multiple (25). It has been reported that SN teeth are more frequently

seen as single, and multiple forms of SN teeth usually occurs as two teeth (15,32). Multiple SN teeth are often found as part of a disease or syndrome.36 In this study, single SN tooth was found to be 92% of the cases; multiple SN teeth were found in 8% of the cases. This finding in the result of our study is consistent with the literature.

Among the main complications created by SN teeth in adjacent teeth and tissues are delayed eruption in adjacent teeth. impaction, midline diastema, loss of vitality, malocclusion, displacement/rotation teeth, adjacent cyst formation, root resorption/dilaceration of adjacent teeth, and eruption towards nasal cavity (35). In our study, at least one complication was found in 40% of cases due to SN teeth. Delayed eruption of adjacent teeth (52.8%) was the most frequently detected complication. Then, it was determined as ectopic tooth eruption (25%), median diastema (13.9%) and malocclusion (8.3%) respectively. The remaining SN teeth (60%) were observed to be asymptomatic. Açıkgöz et al.36 reported that 75% of SN teeth were asymptomatic and most of the cases were diagnosed during routine clinical and radiographic examinations. Demiriz et al.,20, showed that 67.3% of SN teeth were asymptomatic and impaction was the leading complication.

Two individuals who are the carriers of autosomal recessive disorders as a result of consanguineous marriage play a role in the transfer of this gene to the next generation. The rate of consanguineous marriage in Islam societes is quite high. Islam is the dominant religion in Turkey. It was observed that the parents of the children

included study had 75% in our consanguineous marriage. Khan (37) found that there was a relationship between nonsyndromic SN teeth and consanguineous marriage. Studies in Saudi Arabia (38) and Lebanon39 also found that consanguineous marriage was linked to SN teeth. In this study, patients with SN teeth and families were included in an equal number and questioned in terms of consanguineous marriage. Accordingly, the rate consanguineous marriage between parents whose child had SN tooth (76%) and family members who had SN tooth (74%) were found to be similar. We think that such high level of consanguineous marriage might affect the emergence of dental anomalies such as SN teeth.

Today, cone-beam computed tomography (CBCT) is the modern imaging method in dentistry. However. two-dimensional radiograhps are still used in routine dental examination. It is seen that many studies on SN teeth used panoramic radiography archives retrospectively (18,20,21,35). In this study, two-dimensional radiographs (panoramic or intraoral) were used to diagnose SN teeth. CBCT was requested in patients when necessary for surgical operation. In a retrospective study on CBCT, the prevalence of mesiodens type SN teeth was found to be 5.04%. The researchers stated that the use of CBCT in dentistry is advantageous compared to two-dimensional radiographs because it gives detailed information on the subjects such as the location of the SN teeth and their relationship with neighboring anatomical structures (40).

It has been shown that children and adolescents are the most evaluated population for SN teeth in the literature. Most of the patients require surgical tooth extraction before the age of 18, and in some asymptomatic SN teeth patients, followed up for years until they become symptomatic. In this study, it can be seen that complications related to SN teeth may develop in children and adolescents. Careful clinical and radiographic examination in children is important in order to prevent complications that SN teeth may cause, especially in the anterior maxillary region.

## **Acknowledgement:**

The present work was supported by the Research Fund of Istanbul University (Project No. 39623).

## **References:**

- 1. Omer RSM, Anthonappa RP, King NM. Determination of the optimum time for surgical removal of unerupted anterior supernumerary teeth. Pediatr Dent. 2010;32:14–20.
- 2. Pan C-Y, Tseng Y-C, Lan T-H, Chang H-P. Craniofacial features of cleidocranial dysplasia. J Dent Sci. 2017;12:313–8.
- 3. Seehra J, Patel S, Bryant C. Gardner's Syndrome revisited: a clinical case and overview of the literature. J Orthod. 2016;43:59–64.
- 4. Minić S, Novotny GEK, Trpinac D, Obradović M. Clinical features of incontinentia pigmenti with emphasis on oral and dental abnormalities. Clin Oral Investig. 2006;10:343–7.
- 5. Anthonappa RP, King NM, Rabie ABM. Aetiology of supernumerary teeth: a literature review. Eur Arch Paediatr Dent. 2013;14:279–88.

- 6. Sykaras SN. Mesiodens in primary and permanent dentitions: report of a case. Oral Surgery, Oral Med Oral Pathol. 1975;39:870–4.
- 7. Primosch RE. Anterior supernumerary teeth-assessment and surgical intervention in children. Pediatr Dent. 1981;3:204–15.
- 8. Scheiner MA, Sampson WJ. Supernumerary teeth: a review of the literature and four case reports. Aust Dent J. 1997;42:160–5.
- 9. Asaumi J-I, Shibata Y, Yanagi Y, Hisatomi M, Matsuzaki H, Konouchi H, et al. Radiographic examination of mesiodens and their associated complications. Dentomaxillofacial Radiol. 2004;33:125–7.
- 10. Fardi A, Kondylidou-Sidira A, Bachour Z, Parisis N, Tsirlis A. Incidence of impacted and supernumerary teeth-a radiographic study in a North Greek population. Med Oral Patol Oral Cir Bucal. 2011;16:e56-61.
- 11. King N, Tongkoom S, Itthagarun A, Wong H, Lee C. A catalogue of anomalies and traits of the primary dentition of southern Chinese. J Clin Pediatr Dent. 2007;32:139–46.
- 12. Van der Merwe AE, Steyn M. A report on the high incidence of supernumerary teeth in skeletal remains from a 19th century mining community from Kimberley, South Africa: scientific. South African Dent J. 2009;64:162–6.
- 13. Haque S, Alam MK. Common dental anomalies in cleft lip and palate patients. Malaysian J Med Sci MJMS. 2015;22:55.
- 14. Kumar DK, Gopal KS. An epidemiological study on supernumerary teeth: a survey on 5,000 people. J Clin diagnostic Res JCDR. 2013;7:1504.
- 15. Celikoglu M, Kamak H, Oktay H. Prevalence and characteristics of supernumerary teeth in a non-syndrome Turkish population: associated pathologies and proposed treatment. Med Oral Patol Oral Cir Bucal. 2010;15:e575-8.

- 16. Esenlik E, Sayın MÖ, Atilla AO, Özen T, Altun C, Başak F. Supernumerary teeth in a Turkish population. Am J Orthod Dentofac Orthop. 2009;136:848–52.
- 17. Küchler EC, Costa AG da, Costa M de C, Vieira AR, Granjeiro JM. Supernumerary teeth vary depending on gender. Braz Oral Res. 2011;25:76–9.
- 18. Kara Mİ, Aktan AM, Ay S, Bereket C, Şener İ, Bülbül M, et al. Characteristics of 351 supernumerary molar teeth in Turkish population. Med Oral Patol Oral Cir Bucal. 2012;17:e395.
- 19. Kazanci F, Celikoglu M, Miloglu O, Yildirim H, Ceylan I. The frequency and characteristics of mesiodens in a Turkish patient population. Eur J Dent. 2011;5:361.
- 20. Demiriz L, Durmuşlar MC, Mısır AF. Prevalence and characteristics of supernumerary teeth: A survey on 7348 people. J Int Soc Prev Community Dent. 2015;5:S39.
- 21. Karadas M, Celikoglu M, Akdag MS. Evaluation of tooth number anomalies in a subpopulation of the North-East of Turkey. Eur J Dent. 2014;8:337.
- 22. Ferrés-Padró E, Prats-Armengol J, Ferrés-Amat E. A descriptive study of 113 unerupted supernumerary teeth in 79 pediatric patients in Barcelona. Studies. 2009;14:46–52.
- 23. Leco Berrocal M, Martín Morales JF, Martínez González JM. An observational study of the frequency of supernumerary teeth in a population of 2000 patients. Med Oral, Patol Oral y Cirugía Bucal. 2007;12:134–8.
- 24. Fernández Montenegro P, Valmaseda Castellón E, Berini Aytés L, Gay Escoda C. Retrospective study of 145 supernumerary teeth. Med Oral, Patol Oral y Cir Bucal. 2006;11(4):339-344.
- 25. Rajab LD, Hamdan MAM. Supernumerary teeth: review of the literature

and a survey of 152 cases. Int J Paediatr Dent. 2002;12:244–54.

- 26. Zilberman Y, Malron M, Shteyer A. Assessment of 100 children in Jerusalem with supernumerary teeth in the premaxillary region. ASDC J Dent Child. 1992;59:44–7.
- 27. Tay F, Pang A, Yuen S. Unerupted maxillary anterior supernumerary teeth: report of 204 cases. ASDC J Dent Child. 1984;51:289–94.
- 28. Leyland L, Batra P, Wong F, Llewelyn R. A retrospective evaluation of the eruption of impacted permanent incisors after extraction of supernumerary teeth. J Clin Pediatr Dent. 2006;30:225–32.
- 29. Koch H, Schwartz O, Klausen B. Indications for surgical removal of supernumerary teeth in the premaxilla. Int J Oral Maxillofac Surg. 1986;15:273–81.
- 30. DE OLIVEIRA GOMES C, Drummond SN, Jham BC, Abdo EN, Mesquita RA. A survey of 460 supernumerary teeth in Brazilian children and adolescents. Int J Paediatr Dent. 2008;18:98–106.
- 31. Mukhopadhyay S. Mesiodens: a clinical and radiographic study in children. J Indian Soc Pedod Prev Dent. 2011;29:34.
- 32. Sharma A, Singh VP. Supernumerary teeth in Indian children: A survey of 300 cases. Int J Dent. 2012;745265.
- 33. Öztaş B, Bardak Ç, Kurşun EŞ, Akbulut N. Clinical characteristics of non-syndromic

- supernumerary teeth in a cohort of Turkish patients. Oral Radiol. 2011;27:108–13.
- 34. Bello S, Olatunbosun W, Adeoye J, Adebayo A, Ikimi N. Prevalence and presentation of hyperdontia in a non-syndromic, mixed Nigerian population. J Clin Exp Dent. 2019;11:e930.
- 35. Altan H, Akkoc S, Altan A. Radiographic characteristics of mesiodens in a non-syndromic pediatric population in the Black Sea region. J Investig Clin Dent. 2019;10:e12377.
- 36. Acıkgoz A, Acıkgoz G, Tunga U, Otan F. Characteristics and prevalence of non-syndrome multiple supernumerary teeth: a retrospective study. Dentomaxillofacial Radiol. 2006;35:185–90.
- 37. Khan SY. An Exploratory Study of Consanguinity and Dental Developmental Anomalies. Int J Clin Pediatr Dent. 2018;11:513.
- 38. Shokry SM, Alenazy MS. Consanguinity-related hyperdontia: An orthopantomographic study. Dent Res J (Isfahan). 2013;10:732.
- 39. Cassia A, El-Toum S, Feki A, Megarbane A. Five mandibular incisors: an autosomal recessive trait? Br Dent J. 2004;197:307–9.
- 40. Goksel S, Agirgol E, Karabas HC, Ozcan I. Evaluation of Prevalence and Positions of Mesiodens Using Cone-Beam Computed Tomography. J Oral Maxillofac Res. 2018;9

#### **Tables and Charts:**

Table 1: Clinical features of SN teeth.

Variable	Group	Frequency (n)	Percentage (%)
Caralan	Girl	35	35,0
Gender	Boy	65	65,0

	Conical	54	50,0
Т	Tuberculate	18	16,7
Туре	Supplemental	32	29,6
	Odontoma	4	3,7
Name hou	One	92	92,0
Number	Multipl	8	8,0
	Fully impacted	47	42,7
Impaction	Partially erupted	7	6,4
	Erupted	56	50,9
	Vertical	87	80,6
Position	Horizontal	12	11,1
	Inverted	9	8,3
Complication	Yes	40	40,0
Complication	No	60	60,0
	Delayed eruption	38	52,8
True of complication	Malocclusion	6	8,3
Type of complication	Ectopic eruption	18	25,0
	Diastema	10	13,9
Location	Maxilla	108	95,6
Location	Mandible	5	4,4
	Unilateral	63	55,3
Symmetry	Midline	24	21,1
	Bilateral	27	23,7

Table 2: Comparison of SN teeth between dentition groups.

N=108			SN Type					S	
			Conical	Tuberculate	Supplemental	Odontoma	value	d	р
	Mixed	n	19	4	5	1			
Dentiti	dentition	%	65,5%	13,8%	17,2%	3,4%			
on	Permane	n	35	14	27	3	4,312	3	,230
	nt dentition	%	44,3%	17,7%	34,2%	3,8%			

Likelihood Ratio Test

p = 0.05

Table 3: Comparison of SN teeth between gender.

N=108	Type of SN						
	Conical	Tuberculate	Supplemental	Odontoma	value	sd	р

	Girl	n	16	7	13	1			
Gende	GIII	%	43,2%	18,9%	35,1%	2,7%	1 416	2	702
r	Boy	n	38	11	19	3	1,416	3	,702
	БОУ	%	53,5%	15,5%	26,8%	4,2%			

Likelihood Ratio Test

p = 0.05

**Table 4**: Comparison of complication rates between dentition groups.

N=72			Type of complication						
			Delayed eruption	Malocclusion	Ectopic eruption	Diastema	Test value	sd	p
	Mixed	n	9	3	4	4			
Dentiti	dentition	%	45,0%	15,0%	20,0%	20,0%			
on	Permane	n	29	3	14	6	2,631	3	,452
VII.	nt dentition	%	55,8%	5,8%	26,9%	11,5%			

Likelihood Ratio Test

p = 0.05

**Table 5**: The percentage of consanguineous marriage in parents of the cases with SN teeth. Case: SN tooth is seen only in cases, not in family members. Family: SN tooth is detected in at least 1 family member (sibling, mother or father of the child) as well as the child who is examined for SN teeth.

Presence of SN	Case	50	50,0
tooth	Family	50	50,0
Consanguineous	Yes	38	76,0
marriage between parents of cases	No	12	24,0
Concencyingous	Yes	37	74,0
Consanguineous marriage between parents in families	No	13	26,0

Figure 1: Impaction of tooth number 21 due to SN teeth.

