Original Research

Comparing the Incidence of Postdural Puncture Headache (PDPH) between Median and Paramedian Approaches in Nephrolithotripsy under Spinal Anesthesia

Mohammad Sadegh Sanie Jahromi¹, Hasan Zabetian², Navid Kalani³, Fatemeh Khosravani⁴, Alireza Yousefi^{5*}

- 1. Associate Professor of Anesthesiology, Critical care and pain management research center, Jahrom University of Medical Sciences, Jahrom, Iran. Orcid: 0000-0001-8437-1092
- 2. Associate Professor of Anesthesiology, Critical care and pain management research center, Jahrom University of Medical Sciences, Jahrom, Iran. Orcid: 0000-0002-2507-4431
- 3. Anesthesiology, Critical care and pain management research center, Jahrom University of Medical Sciences, Jahrom, Iran. Orcid: 0000-0003-1900-4215
- 4. Student Research Committee, Jahrom University of Medical Sciences, Jahrom, Iran. Orcid: 0009-0004-4486-7113
- 5. Department of Urology, Jahrom University of Medical Sciences, Jahrom, Iran. Orcid: 0000-0002-6415-9227
- *Corresponding Author: Alireza Yousefi, Department of Urology, Jahrom University of Medical Sciences, Jahrom, Iran. Email: a.ryousefi1381@gmail.com

Abstract:

Background: Spinal anesthesia is one of the safest anesthetic procedures with low complications. The most common complication is a headache after surgery or Postdural Puncture Headache (PDPH). It is important to know the factors that effect on PDPH. The aim of this study was to compare the median and paramedian methods in the incidence of PDPH in patients undergoing nephrolithotripsy by spinal anesthesia.

Methods: This clinical trial study was performed on 50 patients who underwent percutaneous nephrolithotomy (PCNL) or nephrolithotripsy referred to Jahrom Peymanieh Hospital in 2019. The patients were randomly divided into two groups of 25 patients. PDPH is defined as bilateral headache in the frontal or occipital region, or neck pain, aggravated by sitting or standing. The data were collected once a day for 7 days after surgery by an uninformed person and then the incidence and prevalence of PDPH were recorded. Patients were followed up by telephone. Data analysis was performed using descriptive and inferential statistics.

Result: No statistically significant distinctions were observed between the two groups in relation to age, gender, and weight. Both the median and paramedian groups exhibited a decrease in the occurrence of postdural puncture headache (PDPH), with a slight elevation noted in the paramedian group. However, no noteworthy variations between the two groups were found on any of the assessed days (p>0.05).

Conclusion: The results of this study showed that there was no difference in the incidence of PDPH incidence in both groups; however, paramedian approach is better than median approach in the frequency of PDPH. Therefore, it is recommended that the paramedian method will be used for spinal anesthesia in patients undergoing PCNL.

Keywords: Paramedian, Median, Spinal anesthesia, PCNL, PDPH

Submitted: 5 Jan 2023, Revised: 27 Jan 2023, Accepted: 9 Feb 2023

Introduction

Subcutaneous nephrolithotomy or nephrolithotripsy (PCNL) is a minimally invasive procedure to remove kidney stones greater than 2 cm (1). In 1976, Fernstorm et al reported for the first time that the removal of calcium kidney stones through a nephrostomy tract (2). Since then, PCNL has been the most common treatment performed for management of renal stones. This technique is less traumatic than open surgery and therefore requires high surgical expertise. There have been many modifications to minimize complications, reduce the need for analgesics, and reduce hospitalization, such as regional block, single-step dilatation, Mini-Perc technique, tubeless PCNL and sandwich therapy (3-5). Nephrolithotripsy can be performed using either general or spinal anesthesia. The utilization of spinal anesthesia offers numerous advantages, including procedural ease, a high success rate, the patient's consciousness during the procedure, reduced postoperative pain, decreased reliance on analgesics, earlier discharge, and avoidance of potential risks associated with general anesthesia and intubation (6-7). Nonetheless, spinal anesthesia carries the potential for certain neurological deficits, such as cauda equina syndrome, hypotension, bradycardia, back pain, shoulder pain, and headaches (7-8). Postdural puncture headache (PDPH) is a major complication of spinal anesthesia that occurs due to inadvertent dural puncture and cerebrospinal fluid leakage and reduced cerebrospinal fluid pressure. Sex, pregnancy, young age, and the common use of neuraxial blocks are risk factors for PDPH (9-10). During epidural anesthesia, inadvertent dural puncture occurs in 1.5% of cases and more than half of these patients develop PDPH (11). Two commonly employed techniques for spinal anesthesia are the median and paramedian approaches, each presenting its own set of advantages and disadvantages. The median

approach, while being the more frequently utilized technique, can pose difficulties, particularly in elderly individuals who may exhibit degenerative changes in the structural components of their spine. On the other hand, the paramedian technique is often favored due to its swifter catheter insertion (12), reduced effort required for needle insertion (13), and the ability to administer anesthesia in an unflexed position (14). Additionally, identifying the epidural space is typically easier with the paramedian technique, and it is associated with fewer osteoarthritic changes elderly patients. However. challenges arise with the oblique position when inserting the epidural needle. It is worth noting that while the paramedian technique reduces the risk of postdural puncture headache (PDPH), its effectiveness has not yet been confirmed in clinical trials (15- 16). Although the use of smaller diameter needles for subarachnoid blocking decreases the likelihood of PDPH, employing such needles for spinal anesthesia can be demanding and result in a lower success rate (17). We typically use spinal needle 25 G in both median and paramedian methods in our hospital. So far, several studies have been performed to compare the median and paramedian methods in terms of PDPH incidence and their results are inconsistent. In the study of Singh B. et al. paramedian technique was significantly preferred technique for reducing PDPH in patients (10). While in Nisar A. et al. study, the incidence of PDPH in the paramedian method was higher than the median, this was not statistically significant (18). Nevertheless, certain studies have reported no significant disparity in the occurrence of postdural puncture headache (PDPH) between the median and paramedian line the approaches (19).In with aforementioned information, our objective was to evaluate and compare the incidence of PDPH in patients undergoing nephrolithotripsy with

spinal anesthesia, specifically focusing on the median and paramedian approaches.

Method

This double-blind randomized clinical trial encompassed the entire patient population referred to Jahrom Peymanieh Hospital in 2019 who underwent percutaneous nephrolithotomy (PCNL). Following approval from the Ethics Committee of the Jahrom University of Medical Sciences, the study aimed to assess and compare the impact of both the median and paramedian approaches to spinal anesthesia on the incidence of postdural puncture headache (PDPH) in patients undergoing nephrolithotripsy. Inclusion criteria included hemodynamic stability (no blood pressure fluctuations), no chronic pain, no use of antihypertensive drugs, no use of anxiolytic and analgesic drugs, no infection at the needle insertion site, no increased Intracranial Pressure (ICP), no coagulation disorders, no hearing problems, and completion of informed consent form for participation in the research. The patients were randomly assigned to one of two groups: 1) Median and 2) Paramedian. Prior to surgery, all patients adhered to the NPO (nil per os) status and received 500 ml of Ringer solution. ECG and pulse oximetry were conducted to monitor blood pressure and heart rate. Measurements of blood pressure and heart rate were recorded at various time points: immediately before spinal anesthesia, following spinal anesthesia, as well as at 5 minutes, 15 minutes, and 30 minutes postspinal anesthesia. After rinsing with Betadine and alcohol, and topical injection of 1 ml of 2% Lidocaine; anesthesia drug (12.5 mg of bupivacaine 0.5%) was injected in sitting position by 25G spinal needle to the subarachnoid space at the L3-4 interspace within 5 seconds, in either median or paramedian method; and immediately after removal of the needle, the patient was in supine position and facial oxygen mask was turned on, 6 liters per minute. According to the

International Headache Association, patients with bilateral headache in the frontal or occipital region or neck pain, which is exacerbated by sitting or standing position and lasts for less than 7 days, were considered as PDPH. PDPH-related data were collected once a day after surgery for 7 days by an uninformed person and the incidence and prevalence of headache were recorded. Patients were followed up by telephone. Inclusion criteria were defined as patients aged 20 to 50 years, interest in study participation, lack of chronic pain and cancer, hemodynamic stability, no anxiety and mental illness, no chronic pain, lack of infection at the needle site, lack of intracranial pressure, and no coagulation disorders. Exclusion criteria also included patients who required medication or other nonroutine care during surgery, and patients who did not meet the inclusion criteria. Data were analyzed using descriptive statistics (mean and standard deviation) and inferential tests (chisquare and t-test) using SPSS software version 21. The level of statistical significance was set at 0.05.

Result

Fifty patients undergoing spinal anesthesia for nephrolithotripsy were divided into two groups: median (25 patients) and paramedian (25 patients). 35 men (70%) and 15 women (30%) were included in the study. The mean age of patients in the median group was 46.88 \pm 12.49 and in the paramedian group 52.48 \pm 12.25 years. There was no significant difference between the two groups. The mean weight of patients in the median group was 77.68 ± 14.29 kg and in the paramedian group was 77.16 ± 9.26 kg. Patients' weight distribution was almost the same in both groups. Percentage of patients with PDPH in median and paramedian groups in the first to seventh days after surgery are showed in the Results showed significant no difference between the two groups in any of the days (P > 0.05), (Table 1).

Changes in systolic blood pressure were compared in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in recovery in table 2. According to the results, systolic blood pressure was significantly higher in the paramedian group before spinal anesthesia, immediately after spinal anesthesia, 5 minutes and 15 minutes after spinal anesthesia (P <0.05) with no significant difference between the two groups at 30 minutes after spinal anesthesia and in recovery (P>0.05).

Patients' diastolic blood pressure changes were compared in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in recovery in table 3.

According to the results, diastolic blood pressure was significantly higher in the paramedian group before the spinal anesthesia, immediately after the spinal anesthesia, 5 minutes and 15 minutes after the spinal anesthesia (p <0.05). No significant difference happened between the two groups at 30 minutes after spinal anesthesia and in recovery (p>0.05).

Patients' heart rate changes were compared in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in recovery in table 4. Results showed that the patients' heart rate was significantly higher in the paramedian group only after 5 minutes of spinal anesthesia (p <0.05). But no significant difference was observed between the two groups in other stages (p> 0.05),(Table 4).

Arterial oxygen saturation changes were compared in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in

recovery in table 5. According to the results, arterial oxygen saturation was significantly higher in the median group before spinal anesthesia, immediately after spinal anesthesia, and 5 minutes after spinal anesthesia (p < 0.05). However, there was no significant difference between the two groups at 15 min and 30 min after spinal anesthesia (p> 0.05). In the recovery, arterial oxygen saturation was significantly higher in the paramedian group (p = 0.047). Arterial oxygen saturation in the median group is decreasing, but this is the ascending in the paramedian group. Although arterial oxygen saturation in paramedian group before spinal anesthesia was higher than paramedian group, this was higher in paramedian group in the recovery phase than median group. (Table 5).

At the end of, the percentage of men and women with PDPH in the median and paramedian groups in the first to seventh days after surgery is showed in table 6 and table 7. According to the results, no significant difference was observed between PDPH rate in both men and women in the median and paramedian groups (p> 0.05). (Table 6).

The study also found that the most frequent use of ephedrine and atropine was in the 5 minutes after spinal anesthesia. In the median group, the most frequent use of ephedrine and atropine was in the 5 minutes after spinal anesthesia. But in the paramedian group, the most frequent use of ephedrine and atropine was in the 15th minute after spinal anesthesia.

In our study, the duration of discharge from recovery was also compared in the median and paramedian groups. There was no significant difference between the two groups in terms of recovery time (p > 0.05). (Table 7).

Discussion

The precise mechanism underlying postdural puncture headache (PDPH) has not yet been fully understood. However, several factors, including the patient's age and gender, can influence its occurrence. PDPH typically arises

from dural leakage of cerebrospinal fluid (CSF) resulting from needle insertion during spinal anesthesia (20-21). In the median approach, the needle traverses the supraspinal and interspinal ligaments, as well as the ligamentum flavum. Conversely, in the paramedian method, the needle directly crosses the ligamentum flavum after passing through the paraspinal muscles, without traversing the supra- and intraspinal ligaments (22). The paramedian approach appears to be simpler than the median approach, particularly in older patients who may have sclerotic ligaments and degenerative changes in the spine and intervertebral spaces (23-24). Singh et al.'s study conducted in India, involving 100 patients undergoing elective surgery, demonstrated no significant difference in the incidence of PDPH between the median and paramedian approaches. They concluded that the paramedian approach significantly reduced the incidence of PDPH, particularly in elderly patients, making it the preferred technique for older individuals Conversely, a study by Nisar et al. in 2016, comprising 100 patients in Pakistan, reported a significantly higher incidence of PDPH in the paramedian technique compared to the median technique (18). In our study, the incidence of PDPH decreased in both the median and paramedian groups, with a slight increase observed in the paramedian group, although no significant difference was found between the two groups. These findings are consistent with the results obtained by Firdous et al. Their study also suggested that the paramedian approach was superior to the median approach in reducing PDPH, albeit without statistical significance (26). In our study, there was no significant difference between the two groups in terms of age, weight, and sex, which aligns with the findings of Singh et al.'s study (25). As mentioned earlier, gender is considered a risk factor for PDPH. Some studies have reported no significant difference in the incidence of PDPH between men and women (20, 27-28),

which corresponds with our study's results. However, a study conducted by DelPizzo et al. in 2017, involving 300 patients undergoing knee surgery in New York, indicated a higher incidence of PDPH in women compared to men (29). Our study revealed that systolic and diastolic blood pressure were significantly higher in the paramedian group before anesthesia, immediately after anesthesia, as well as at 5 minutes and 15 minutes after anesthesia. Nonetheless, Chong et al.'s study that PDPH was unaffected found hypertension (30). In a study by Kim et al. in 2013, the results demonstrated a decrease in blood pressure and heart rate following spinal anesthesia (31), which is consistent with our study's findings concerning systolic and diastolic blood pressure. However, in our study, the heart rate increased 15 minutes after anesthesia compared to before anesthesia but then decreased from 30 minutes after anesthesia in both the median and paramedian groups.

Conclusion

The results of this study showed that there was no difference in the incidence of PDPH incidence in both groups; however, paramedian approach is better than median approach in the frequency of PDPH. Therefore, it is recommended that the paramedian method will be used for spinal anesthesia in patients undergoing PCNL.

Ethical Code

This project has been approved by the research ethic committee, as well as, supported with grant number of "IR.JUMS.REC.1397.132" by Jahrom University of Medical Sciences in Jahrom, Iran.

References

1. Malik I, Wadhwa R. Percutaneous Nephrolithotomy: Current Clinical Opinions and Anesthesiologists Perspective. Anesthesiology Research and Practice. 2016;2016:7.

- 2. Fernström I, Johansson B. Percutaneous Pyelolithotomy. Scandinavian Journal of Urology and Nephrology. 1976 1976/01/01;10(3):257-9.
- 3. Goh M, Wolf JS, Jr. Almost totally tubeless percutaneous nephrolithotomy: further evolution of the technique. Journal of endourology. 1999 Apr;13(3):177-80.
- 4. Jackman SV, Hedican SP, Peters CA, Docimo SG. Percutaneous nephrolithotomy in infants and preschool age children: experience with a new technique. Urology. 1998;52(4):697-701.
- 5. Saied MM, Sonbul ZM, el-Kenawy M, Atallah MM. Spinal and interpleural bupivacaine for percutaneous nephrolithotomy. Middle East journal of anaesthesiology. 1991 Oct;11(3):259-64.
- 6. Asgari Z, Rezaeinejad M, Hosseini R, Nataj M, Razavi M, Sepidarkish M. Spinal Anesthesia and Spinal Anesthesia with Subdiaphragmatic Lidocaine in Shoulder Pain Reduction for Gynecological Laparoscopic Surgery: A Randomized Clinical Trial. Pain research & management. 2017:2017:1721460.
- 7. Ortiz J, Rajagopalan S. A Review of Local Anesthetic Techniques for Analgesia After Laparoscopic Surgery. J Minim Invasive Surg Sci. 2014;3(2):e11310.
- 8. Jun G-W, Kim M-S, Yang H-J, Sung T-Y, Park D-H, Cho C-K, et al. Laparoscopic appendectomy under spinal anesthesia with dexmedetomidine infusion. Korean journal of anesthesiology. 2014;67(4):246-51. PubMed PMID: 25368782.
- 9. Kwak K-H. Postdural puncture headache. Korean Journal of Anesthesiology. 2017 04/01;70:136.
- 10. Singh B, Sohal A, Singh I, Goyal S, Kaur P, Attri J. Incidence of postspinal headache and low backache following the median and paramedian approaches in spinal anesthesia. Anesthesia: Essays and

- Researches. 2018 January 1, 2018;12(1):186-9.
- 11. Choi PT, Galinski SE, Takeuchi L, Lucas S, Tamayo C, Jadad AR. PDPH is a common complication of neuraxial blockade in parturients: a meta-analysis of obstetrical studies. Canadian journal of anaesthesia = Journal canadien d'anesthesie. 2003 May;50(5):460-9.
- 12. Leeda M, Stienstra R, Arbous MS, Dahan A, Th Veering B, Burm AG, et al. Lumbar epidural catheter insertion: the midline vs. the paramedian approach. European journal of anaesthesiology. 2005 Nov;22(11):839-42.
- 13. Blomberg RG, Jaanivald A, Walther S. Advantages of the paramedian approach for lumbar epidural analgesia with catheter technique. A clinical comparison between midline and paramedian approaches. Anaesthesia. 1989 Sep;44(9):742-6.
- 14. Podder S, Kumar N, Yaddanapudi LN, Chari P. Paramedian lumbar epidural catheter insertion with patients in the sitting position is equally successful in the flexed and unflexed spine. Anesthesia and analgesia. 2004 Dec;99(6):1829-32
- 15. Janik R, Dick W. [Post spinal headache. Its incidence following the median and paramedian techniques]. Der Anaesthesist. 1992 Mar;41(3):137-41.
- Mosaffa F, Karimi K, Madadi F, Khoshnevis SH, Daftari Besheli L, Eajazi A. Post-dural Puncture Headache: A Comparison Median Between and Paramedian Approaches in Orthopedic Patients. Anesth Pain Med. 2011 Fall;1(2):66-9. PubMed PMID: 25729658.
- 17. Ghaleb A. Postdural puncture headache. Anesthesiol Res Pract. 2010;2010. PubMed PMID: 20814596. Pubmed Central PMCID: PMC2931374.
- 18. Nisar A, Saleem J, Hussain S, Bashir K. Comparison of postduralpuncture headache in median and paramedian approach under

- spinal anesthesia in cesarean section. Pak J Med Health Sci 2016;10(1):298-301.
- 19. Bapat V, Vishwasrao S. Spinal anaesthesia with midline and paramedian technique in elderly patients. Indian J Appl Res 2015;5(9):442-4.
- 20. Dittmann M, Schaefer HG, Renkl F, Greve I. Spinal anaesthesia with 29 gauge Quincke point needles and post dural puncture headache in 2,378 patients. Acta anaesthesiologica Scandinavica. 1994 Oct;38(7):691-3. PubMed PMID: 7839778.
- 21. Reid JA, Thorburn J. Headache after spinal anaesthesia. British journal of anaesthesia. 1991 Dec;67(6):674-7. PubMed PMID: 1768536.
- 22. Haider S, Butt KJ, Aziz M, Qasim M. Postdural puncture headache A comparison of midline and paramedian approaches. Biomedica. 2005;21:90-2.
- 23. Ahmed T, Ullah H. Paramedian technique of spinal anaesthesia in elderly patients for hip fracture surgery. Journal of the College of Physicians and Surgeons-Pakistan: JCPSP. 2007 Mar;17(3):184.
- 24. Ahsan-ul-Haq M, Amin S, Javaid S. Paramedian technique of spinal anesthesia in elderly patients for hip fracture surgery. Journal of the College of Physicians and Surgeons--Pakistan: JCPSP. 2005 Mar;15(3):160-1.25. Singh P, Agrawal SK, Dwivedi S, Yadav A. A comparative evaluation between median and paramedian approaches for sub-

- arachnoid block in elderly patients. 2017. 2017 2017-01-02;4(6):4.
- 26. Firdous T, Siddiqui MA, Siddiqui SM. Frequency of post dural puncture headache in patients undergoing elective cesarean section under spinal anesthesia with median versus paramedian approach. Anaesthesia, Pain and Intensive Care. 2016 01/01;20:165-70.
- 27. Lybecker H, Moller JT, May O, Nielsen HK. Incidence and prediction of postdural puncture headache. A prospective study of 1021 spinal anesthesias. Anesthesia and analgesia. 1990 Apr;70(4):389-94.
- 28. Tarkkila PJ, Heine H, Tervo RR. Comparison of Sprotte and Quincke needles with respect to post dural puncture headache and backache. Regional anesthesia. 1992 Sep-Oct;17(5):283-7.
- 29. DelPizzo K, Cheng J, Dong N, Edmonds CR, Kahn RL, Fields KG, et al. Post-Dural Puncture Headache is Uncommon in Young Ambulatory Surgery Patients. HSS J. 2017;13(2):146-51.
- 30. Chung JM, Kim JK, Kim JY, Park KI, Yu HA. Post-dural puncture headache after spinal anesthesia and perioperative blood pressure. Journal of the Neurological Sciences. 2013;333:e504-e5.
- 31. Kim HY, Lee MJ, Kim M-N, Kim J-S, Lee WS, Lee KC. Effect of position changes after spinal anesthesia with low-dose bupivacaine in elderly patients: sensory block characteristics and hemodynamic changes. Korean journal of anesthesiology. 2013;64(3):234-9.

Table 1: The incidence of PDPH in patients in the median and paramedian groups in the first to seventh days

Group	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Median	42.1%	40%	25.0%	10.0%	10.0%	5.0%	5.0%
Paramedian	50%	50%	14.3%	0.0%	0.0%	0.0%	0.0%
P value	.923	.820	.742	.632	.632	1.000	1.000

Table 2: Systolic blood pressure changes in median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in recovery

Group		before	after	5 minutes	15 minutes	30 minutes	recovery
Median	Mean	135.00	128.84	116.20	117.24	118.72	118.17
1.1culuii	SD	21.89	28.18	21.17	16.91	18.04	18.28
Paramedian	Mean	148.24	144.52	133.44	133.00	115.00	123.73
	SD	18.66	15.43	18.96	19.44	17.53	12.95
Pv	alue	.026	.018	.004	.004	.463	.248

Table 3: Diastolic blood pressure changes in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes, and 30 minutes after spinal anesthesia and in recovery

Group		Before	After	5 minutes	15 minutes	30 minutes	recovery
Median Mean SD	Mean	87.12	79.84	72.80	76.04	75.88	72.96
	SD	15.64	18.27	13.22	11.57	14.84	12.74
Paramedian	Mean	95.48	90.08	86.48	149.48	75.44	72.36
Taramedian	SD	7.67	8.45	14.88	173.17	12.74	10.36
P value		.020	.014	.001	.040	.911	.865

[Downloaded from intjmi.com on 2025-10-30]

Table 4: Heart rate changes in the median and paramedian groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes and 30 minutes after spinal anesthesia and in recovery

Group		Before	After	5 minutes	15 minutes	30 minutes	recovery
Median	Mean	83.52	91.20	80.56	98.08	85.76	78.30
	SD	11.20	19.60	14.62	1.85	17.42	15.72
Paramedian	Mean	88.08	94.04	99.12	97.24	85.68	81.82
	SD	12.58	17.01	17.48	2.37	13.41	14.04
P value		.182	.587	<0.001	.168	.986	.434

Table 5: Arterial oxygen saturation changes in the median and paramedine groups before spinal anesthesia, immediately after spinal anesthesia, 5 minutes, 15 minutes and 30 minutes after spinal anesthesia and in recovery

Group		Before	After	5 minutes	15 minutes	30 minutes	recovery
Median .	Mean	98.88	98.80	98.60	98.08	98.40	96.17
	SD	1.27	1.19	1.47	1.85	1.55	7.20
Paramedian	Mean	96.60	97.96	97.28	97.24	97.88	100.18
	SD	2.57	1.31	1.95	2.37	4.36	5.85
P value		< 0.001	.022	.009	.168	.577	.047

Table 6: The incidence of PDPH in median group on day 1 to day 7 by sex

Group	Sex	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Median	Male	38.5%	42.9%	28.6%	7.1%	7.1%	7.1%	7.1%
177CGTGTT	Female	50.0%	33.3%	16.7%	16.7%	16.7%	0.0%	0.0%
P value		1.000	1.000	1.000	1.000	1.000	1.000	1.000

Table 7: The incidence of PDPH in paramedian group on day 1 to day 7 by sex

Group	Sex	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Median	Male	63.6%	63.6%	18.2%	0.0%	0.0%	0.0%	0.0%
Wicdian	Female	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
P value		.193	.193	1.000				